

CORESTA 2024

Presentation no. ST16

Influence of nicotine concentration and flavours on mouth level exposure and puffing topography among regular e-cigarette consumers in New Zealand

Lauren Edward, Krishna Prasad, Linsey Haswell, Adam Gray, Tahseen Jilani, Stacy Fiebelkorn



B.A.T. (Investments) Limited, Regents Park Road, Millbrook, Southampton SO15 8TL, UK

This presentation is the property of BAT. Exclusively for discussion purposes.



FOR DISCUSSION PURPOSES ONLY

PRESENTATIONS ARE INTENDED FOR INFORMATIONAL PURPOSES ONLY. STATEMENTS OF FACT AND OPINIONS EXPRESSED ARE THOSE OF THE PRESENTER ONLY, AND ARE NOT NECESSARILY THE OPINIONS, POSITIONS OR POLICIES OF BRITISH AMERICAN TOBACCO (INVESTMENTS) LIMITED OR ITS AFFILIATES (“BAT”). BAT DOES NOT ENDORSE OR APPROVE, AND ASSUMES NO RESPONSIBILITY FOR THE CONTENT, ACCURACY OR COMPLETENESS OF THE INFORMATION PRESENTED. THESE MATERIALS MAY PRESENT CONCEPTS, EXECUTIONS, IDEAS AND THEMES THAT ARE NOT LEGALLY PERMISSIBLE (EITHER IN WHOLE OR IN PART) OR ACCEPTABLE IN CERTAIN JURISDICTIONS, OR WHICH MAY INFRINGE THIRD-PARTY RIGHTS. THIS PRESENTATION MAY CONTAIN REFERENCES TO LAWS AND REGULATIONS THAT MAY CHANGE OVER TIME AND SHOULD BE INTERPRETED IN THAT CONTEXT.

THIS PRESENTATION IS INTENDED FOR SCIENTISTS AND POLICYMAKERS. WE ENCOURAGE PERSONS INTERESTED IN TOBACCO HARM REDUCTION TO CONSIDER ALL POINTS OF VIEW AND SOURCES OF INFORMATION. BAT DOES NOT MAKE HEALTH CLAIMS REGARDING ITS BRANDS. NOTHING CONTAINED HERE SHOULD BE MISCONSTRUED TO THE CONTRARY. TO THE EXTENT THAT THIRD-PARTY SOURCES ARE REFERENCED, NEITHER THE PRESENTER NOR BAT IS RESPONSIBLE FOR THE CONTENT OF REFERENCED SOURCES AND THE VIEWS EXPRESSED MAY NOT REPRESENT THE VIEWS OF THE PRESENTER OR BAT. NO TOBACCO PRODUCT IS SAFE, ALL TOBACCO PRODUCTS CONTAINING NICOTINE ARE ADDICTIVE. YOUTH SHOULD NEVER USE TOBACCO. SMOKERS WHO ARE CONCERNED ABOUT THEIR HEALTH SHOULD QUIT.

THE TOPICS, CONCEPTS AND OTHER INFORMATION DISCUSSED HEREIN ARE NOT FINAL, ARE SUBJECT TO CHANGE AND/OR CANCELLATION AND MAY BE FOR ILLUSTRATIVE OR THEORETICAL PURPOSES ONLY. NO DEFINITIVE PLANS OR COMMITMENTS SHOULD BE INFERRED FROM THESE MATERIALS AND ANY PROPOSED PLANS OR COMMITMENTS ARE SUBJECT IN ALL RESPECTS TO APPLICABLE INTERNAL REVIEW AND GOVERNANCE REQUIREMENTS AND LAWS AND REGULATIONS IN RESPECTIVE JURISDICTIONS.

Contents

- 1 Background & objectives
- 2 Study design & methodology
- 3 Results: influence of nicotine concentration
- 4 Results: influence of flavour
- 5 Results: nicotine delivery comparison with a tobacco cigarette
- 6 Conclusions



Background



Vapour product aerosols have a reduced toxicant profile relative to cigarette smoke [1-3]



Consumer's actual exposure is influenced by how they use the product (e.g., puff duration and use frequency)



Use behaviour may be influenced by device characteristics (e.g., power, open pressure drop, e-liquid flavour and nicotine concentration)



Many studies on the impact of nicotine concentration on product use have been small scale, low participant numbers and few conditions (high vs low nicotine)

[1] Shah, N.H., et al., Non-targeted analysis using gas chromatography-mass spectrometry for evaluation of chemical composition of E-vapor products. *Front Chem*, 2021. 9:742854. DOI: 10.3389/fchem.2021.742854
[2] Wagner, K.A., et al., An evaluation of electronic cigarette formulations and aerosols for harmful and potentially harmful constituents (HPHCs) typically derived from combustion. *Regul Toxicol Pharmacol*, 2018. 95: p. 153-160. DOI: 10.1016/j.yrtph.2018.03.012
[2] Rudd, K., et al., Chemical composition and in vitro toxicity profile of a pod-based e-cigarette aerosol compared to cigarette smoke. *Appl In Vitro Toxicol*, 2020. 6(1): p. 11-41. DOI: 10.1089/aivt.2019.0015



Objectives



Primary Objective

Measure the puffing topography, mouth level exposure (MLE) to aerosol and nicotine, and average daily consumption (ADC) among vapour product consumers when using a closed rechargeable system with e-liquids of five nicotine concentrations (6 – 47 mg/mL) and four flavours



Secondary Objective

Measure the puffing topography & MLE of smokers when using a cigarette for comparison



Vuse ePod 2

Vapour products

Vuse ePod 2	E-liquid nicotine level (mg/mL)				
	6	12	18	34	47
Crisp Mint	X	X	X	X	X
Golden Tobacco			X		
Tropical Mango			X		
Peppermint Tobacco			X		

Cigarette


Rothmans Royal	15 mg ISO tar
-----------------------	----------------------

Study Design



Central Location Test (Christchurch, New Zealand): Puffing topography & MLE via desktop puffing analyser; sensory perception via questionnaire

Home Use Test: Puffing topography via connected product use behaviour (PUB) device; ADC and sensory perception via questionnaire



Methodology




Central Location Testing

- Desktop puffing analyser (PA1) – puff volume, duration, interpuff interval (IPI), flow rate, optical obscuration of aerosol, number of puffs measured in real-time ^[1-2]
- Participants used the product through the PA1 holder
- MLE to aerosol and nicotine for vapour product estimated from DML by vaping machine calibration ^[2]
- MLE to NFDPM and nicotine for cigarette estimated from per-puff optical obscuration by machine calibration ^[1]

Home Use Testing

- Connected *product use behaviour* (PUB) device - duration, IPI, angle held, date, time and number of puffs ^[3]
- Connects between battery section and e-liquid pod
- MLE to aerosol and nicotine estimated from puff duration by vaping machine calibration



Abbreviations: MLE = mouth level exposure; NFDPM = nicotine-free dry particulate matter

1-Slayford S, Frost B. A device to measure a smokers' puffing topography and real-time puff by puff "tar" delivery. *Beit Tab Int.* 2014;26(18):74-84

2-Jones J et al., A cross-category puffing topography, mouth level exposure and consumption study among Italian users of tobacco and nicotine products. *Sci Rep.* 2020;10(1):12

3-Underly RG et al., Dull GM, Nudi E, Pionk T, Prevette K, Smith J (2023). Using a Novel Connected Device for the Collection of Puffing Topography Data for the Vuse Solo Electronic Nicotine Delivery System in a Real-World Setting: Prospective Ambulatory Clinical Study. *JMIR Form Res.* 7: e49876. doi:10.2196/49876

This presentation is the property of BAT. Exclusively for discussion purposes.



RESULTS


Effect of nicotine concentration

Central Location Testing*


With increasing concentration of nicotine in e-liquid:

Overall trends

- Mean puff duration decreased
- IPI increased (reduced puff frequency, i.e. fewer puffs per 15-min)



- Estimated MLE to nicotine (per 15-min session & per puff) increased
- MLE to aerosol per session decreased



Abbreviations: MLE = mouth level exposure; IPI = inter-puff interval

* Data are mean from 90 participants who used each product twice. Error bars indicate standard error. Data were analysed by mixed-effects ANOVA, followed by Tukey's post-hoc test. For a given parameter, values sharing the same alphabet letter were not significantly different ($p > 0.05$); those not sharing the same alphabet letter were significantly different ($p < 0.05$). Participants were given a maximum of 15-min in which to use the product.


This presentation is the property of BAT. Exclusively for discussion purposes.

Central Location Testing*


With increasing concentration of nicotine in e-liquid:

Trends across nicotine user groups

- 18 mg/mL users took fewer puffs than 34 & 47 mg/mL users
- No differences in other topography attributes among user groups



- At higher nicotine concentrations – lower estimated MLE to nicotine for regular 18 mg/mL users compared with 34 & 47 mg/mL users



Abbreviations: MLE = mouth level exposure

* Participants used each product twice. Error bars indicate standard error. Participants were given a maximum of 15-min in which to use the product.


This presentation is the property of BAT. Exclusively for discussion purposes.

Central Location Testing*

With increasing concentration of nicotine in e-liquid:

Overall trends

- Perceived intensity and irritation increased
- No differences in overall liking or other sensory perception attributes



Abbreviations: MLE = mouth level exposure

* Data are mean from 90 participants who used each product twice. Error bars indicate standard error. Data were analysed by mixed-effects ANOVA, followed by Tukey's post-hoc test. For a given parameter, values sharing the same alphabet letter were not significantly different ($p > 0.05$); those not sharing the same alphabet letter were significantly different ($p < 0.05$). Participants were given a maximum of 15-min in which to use the product.



This presentation is the property of BAT. Exclusively for discussion purposes.

Home Use Testing*

When participants used their usual nicotine concentration:

- 47 mg/mL users puffed less frequently than 18 and 34 mg/mL users
- No differences in puff duration (1.58-1.67 s) or ADC (0.6-0.75 pods/day)

- No significant differences in estimated MLE per day
- However, MLE to nicotine per day was lowest for 18 mg/mL users



Abbreviations: ADC = average daily consumption (self-reported via daily consumption diary); MLE = mouth level exposure.

* Participants used only their usual nicotine concentration (either 18, 34 or 47 mg/mL). Only participants with valid data are included. Error bars indicate standard error. Data were analysed by one-way ANOVA, followed by Tukey's post-hoc test. For a given parameter, values sharing the same alphabet letter were not significantly different ($p > 0.05$); those not sharing the same alphabet letter were significantly different ($p < 0.05$). Participants were given a maximum of 15-min in which to use the product.

This presentation is the property of BAT. Exclusively for discussion purposes.





RESULTS

Effect of flavour liking

Central Location Testing*

Across flavours at 18 mg/mL nicotine:

- Shorter puff duration, fewer puffs, and lower MLE for Golden Tobacco flavour, consistent with lower taste liking rating (1.8 vs 3.2 -3.8**) and lower overall liking rating (2.0 vs 3.3 – 3.7**)
- Slightly larger puffs of longer duration when using Mango flavour (55 vs 46 – 50 mL puff volume; 2.6 vs 2.1 – 2,3 s puff duration)



Abbreviations: MLE, = mouth level exposure

* Data are mean from 90 participants who used each product twice. Error bars indicate standard error. Data were analysed by mixed-effects ANOVA, followed by Tukey's post-hoc test. For a given parameter, values sharing the same alphabet letter were not significantly different ($p > 0.05$); those not sharing the same alphabet letter were significantly different ($p < 0.05$). Participants were given a maximum of 15-min in which to use the product.



** Via self-reported questionnaire, scored on a scale of 1-5 where 1 = "dislike a lot" and 5 = "like a lot"

This presentation is the property of BAT. Exclusively for discussion purposes.

Home Use Testing*

Across flavours at 18 mg/mL nicotine:

- Fewer puffs and shorter puff duration for Golden Tobacco flavour consistent with findings in CLT
- Higher number of puffs for Crisp Mint flavour (not statistically significant) may be attributable to accessibility of this flavour in New Zealand
- No significant differences in MLE to aerosol or nicotine



Abbreviations: MLE = mouth level exposure; CLT = central location test

* Only participants with valid data are included. Error bars indicate standard error. Data were analysed by mixed-effects ANOVA, followed by Tukey's post-hoc test. For a given parameter, values sharing the same alphabet letter were not significantly different ($p > 0.05$); those not sharing the same alphabet letter were significantly different ($p < 0.05$). Participants were given a maximum of 15-min in which to use the product.

This presentation is the property of BAT. Exclusively for discussion purposes.





RESULTS

Nicotine delivery -
comparison with a tobacco cigarette

Nicotine delivery: vapour vs cigarette*

Central location testing - estimated MLE to nicotine: vapour product vs cigarette

- Estimated MLE to nicotine per puff for 47 mg/mL vapour product was significantly lower than 15 mg cigarette
- Vapers took significantly more puffs per 15-min vaping session than cigarette smokers took per cigarette
- Thus, total estimated MLE to nicotine for 47 mg/mL vapour product was similar to the cigarette



Abbreviations: MLE = mouth level exposure

* Data are mean from 90 e-cigarette users (using only the e-cigarette) and 35 cigarette smokers (using only the cigarette) who used each product twice. Error bars indicate standard error. Data were analysed by mixed-effects ANOVA, followed by Dunnett's post-hoc test. For a given parameter, values that do not share the letter A are significantly different from the cigarette ($p < 0.05$).

This presentation is the property of BAT. Exclusively for discussion purposes.

Conclusions



Puffing behaviour was broadly consistent with ISO20768:2018 machine puffing regime

- Average puff durations (1.5 -2.6s) were less than the 3s recommended by CORESTA
- Puff volumes (46 – 55mL) and IPI (35 – 143s) were broadly consistent with the recommended 55mL volume and 30s frequency



Both nicotine level and flavour influence consumers' puffing behaviour

- Puff duration and puff frequency tended to decrease with increasing nicotine concentration and this may be associated with self-titration or higher perceived irritation
- Flavours with a lower overall liking score tended to result in reduced use behaviour



Nicotine concentration and e-liquid flavour play an important role in helping adult smokers who would otherwise continue to smoke to instead switch to Vapour Products



Thank you for listening

Any questions?



Contact: Lauren_Edward@bat.com

